

IN VITRO EMBRYOGENESIS OF *NEOMYSIS INTEGER* (CRUSTACEA, MYSIDACEA) AS A POTENTIAL INDICATOR OF ENDOCRINE DISRUPTION

Fockedey Nancy¹, Stijn Bruwiere¹, An Ghekiere² and Magda Vincx¹

¹ Marine Biology Section, Departement of Biology, Ghent University
Krijgslaan 281/S8, B-9000 Ghent, Belgium
E-mail: nancy.fockedey@ugent.be

² Laboratory of Environmental Toxicology and Aquatic Ecology, Ghent University
J. Plateastraat 22, B-9000 Ghent, Belgium

The brackish water mysid *Neomysis integer* has been put forward to use as a test organisms for the evaluation of environmental endocrine disruption in the brackish reaches of Western European estuaries and inland water bodies. As the embryonic development is a critical time window within the life history of an organism, it has high potential to serve as a bio-assay for testing endocrine disruption. Within the present study a protocol has been developed to follow the in vitro embryogenesis of *Neomysis integer* and the most sensitive endpoints for endocrine disruption has been determined.

Daily survival percentage, percentage survival days, hatching success, duration of each developmental stage and the size increment of the embryos were followed as potential endpoints. The optimal combination of salinity and temperature was obtained using a central composite design, and the response of the endpoints was fitted with a response surface model. The survival and hatching success are highly dependent on the salinity conditions, while the development time is strongly affected by temperature. Optimal salinity for embryonic development of *Neomysis integer* is 15psu; optimal temperature around 15 - 20°C.

The potential of the endpoints for use as a bio-assay for endocrine disruption was tested with the pesticide methopreen, a juvenile hormone-analog. Post-fertilization exposure of the embryos at 1µg to 100µg methopreen l⁻¹ caused a significantly lower survival and hatching percentage. As a result of concentrations from 0.01µg methopreen l⁻¹ and higher, the development time of the embryonic stage II and III and the time to hatching were altered.